

# STOICHIOMETRY


Measurements in Chemical  
Reactions

# STOICHIOMETRY

- The analysis of the quantities of substances in a chemical reaction.
- Stoichiometric calculations depend on the MOLE-MOLE relationships of substances.
- Therefore, a BALANCED chemical equation is required.

# Mole-Mole Relationships

- Consider the acid-base neutralization:




Number of moles BEFORE Reaction	5	? =	0	0
Number of moles AFTER Reaction	? =	? =	? =	? =

- How many moles of sodium hydroxide would be needed to completely react with 5 moles of sulfuric acid?
- How many moles of water and sodium sulfate would be formed?

# Mole-Mole Relationships

- Consider the acid-base neutralization:




Number of moles BEFORE Reaction	? =	6	0	0
Number of moles AFTER Reaction	? =	? =	? =	? =

- How many moles of magnesium hydroxide would be needed to completely react with 6 moles of phosphoric acid?
- How many moles of water and magnesium phosphate would be formed.

# Mass-Mass Problem

- We can convert from the mass of a reactant or product to the mass of any other reactant or product.
- However, to do this, we must always convert from one substance to the other **through moles**.

- **EXAMPLE:** Consider the combustion of 8.96 g hexane:



- *What mass of oxygen will be required for the complete combustion of the hexane?*
- *What masses of carbon dioxide & water will be formed by this reaction?*

## Mass – Mass Problem (Continued)

- It is often helpful to set up a table to organize information that is needed.




molar mass (g/mol)	86.18	32.00	44.01	18.02
mass (g)	8.96			
Mole				

## General Stoichiometry Steps:

1. Convert quantity of given substance to moles.
2. Convert moles of *given* substance to moles of the *unknown* substance.
3. Convert moles of the unknown substance to the desired units.

## Stoichiometry Example

- In the reaction below, what mass of lead (II) chloride can be produced from the reaction of 75.0 g of aluminum chloride with excess lead (II) nitrate?




molar mass	133.33 g/mol			278.11 g/mol
mass	75.0 g	<i>Excess (more than needed)</i>		
mole				



## Stoichiometry Example

What mass of chlorine gas is required to fully react with excess aluminum powder to produce 2.50 kg aluminum chloride?



Yield: quantity of product(s).

- **Theoretical Yield:** the calculated amount of product that can be formed in a chemical reaction if ALL of the reactants fully react. (Usually expressed in grams.)
- **Actual Yield:** the amount of product actually produced in a laboratory experiment.
- **Percent Yield** =  $\frac{\text{Actual}}{\text{Theoretical}} \times 100\%$

## Theoretical Yield

- An excess of elemental iodine is reacted with 1.20 g of  $\text{NaBH}_4$  to obtain 0.295 g of  $\text{B}_2\text{H}_6$ . What is the percent yield of  $\text{B}_2\text{H}_6$ ?



- 0.295 g  $\text{B}_2\text{H}_6$  is the ACTUAL yield. It is NOT part of the stoichiometry problem
- Hint: Find the theoretical yield first.

# Limiting Reagents & Oreo Cookies



2 cookies + 1 crèmes → 1 OREO

# Oreo Cookies

How many OREOS can we make if we have 10 cookies and 5 crèmes?

**2 cookies + 1 crèmes → 1 OREO**

Initial	10	5	0
Change	- 10	- 5	+ 5
Final	0	0	5

We make 5 OREO's with nothing left over.

# Oreo Cookies

How many OREOS can we make if we have 20 cookies and 15 crèmes?



Initial	20	15	0
Change			
Final			

In this case, the cookies are the *limiting reagent*, and the crèmes are *in excess*.

# Oreo Cookies

How many OREOS can we make if we have 14 cookies and 6 crèmes?

**2 cookies + 1 crèmes → 1 OREO**

Initial	14	6	0
Change			
Final			

What is the limiting ingredient?

What will be left over?


# Definitions

- The ***LIMITING REAGENT*** (or limiting reactant) is the reactant that runs out first in a chemical reaction.
- The ***EXCESS REAGENT*** is the reactant present in a quantity greater than needed to fully react with the L.R.






- When 10 moles of hydrogen is mixed with 10 moles of oxygen and allowed to react, 10 moles of water is produced, with 5 moles oxygen left over.



## Limiting Reagent Example


- If 5 mol of nitrogen gas and 12 mol of hydrogen gas are mixed and allowed to react to form ammonia,
  - What is the limiting reagent?
  - How many moles of  $\text{NH}_3$  can be formed?
  - How many moles of the excess reagent will be left over?



Initial			
Change			
Final			


## Example – Limiting Reagents

Nitrogen gas can be reacted with sodium metal at high pressure to make sodium azide. What is the limiting reagent if 5.50 g of sodium is mixed with 8.00 g of nitrogen?



## Example – Limiting Reagents

Consider the reaction of 5.00 g of elemental sulfur,  $S_8$ , with 5.00 g of chlorine gas,  $Cl_2$ , to form  $SCl_3(l)$ .



- A) What is the Limiting Reagent?
- B) What is the theoretical yield (in g) of sulfur trichloride?
- C) What mass of the excess reagent will remain?

## A) What is the Limiting Reagent?

- We must compare moles to moles, taking into account the **STOICHIOMETRY** (mole ratios) of the reaction.
- One method to determine which reactant will provide fewer moles of the desired product.
- The one that could produce less product is the **LIMITING REAGENT**.

## B) What mass of sulfur trichloride will be produced?

- Use the number of moles of the product that the limiting reagent will produce to determine the mass of product that is expected.
- The *limiting reagent* **always** determines what the *theoretical yield* of the products will be.

C) What mass of the excess reagent will remain?

Determine how much of the excess reagent is required to *react with* the given quantity of the limiting reagent.

*Mass remaining =*

*Initial mass – Mass reacted*

# Stoichiometry Example

Consider the preparation of titanium metal:



If 2.55 kg of titanium (IV) chloride is reacted with 1.00 kg of magnesium metal,

- A. What is the limiting reagent?
- B. What is the theoretical yield of titanium?
- C. What mass of excess reagent remains?



## Other routes to the MOLE (starting from volumes)

- **Density ( $\text{g}/\text{cm}^3$ )** – allows you to convert the volume of a solid or a pure liquid (not a solution) to grams, which may then easily be converted to moles using molar mass.
- **Molar Gas Density ( $22.4 \text{ L}/\text{mol}$ )** – One mole of any IDEAL gas will occupy 22.4 L at 1.00 atm pressure and  $0^\circ\text{C}$  (also known as STP conditions – standard temperature and pressure.) This conversion factor can be used to convert volume of a gas to moles.
- **Concentration in Molarity ( $\text{mol}/\text{L}$ )** – Molar concentrations can be used to convert directly from the volume of a solution (easy to measure in the lab) to moles of the solute.

## Stoichiometry Example

Consider the reaction:  $2 \text{K}_{(s)} + \text{I}_{2(s)} \rightarrow 2 \text{KI}_{(s)}$

A chemistry lab student began with 125.0 g of potassium and 40.5 cm<sup>3</sup> of iodine. When the product was obtained and massed, the student found that 244.2 g of KI had actually been produced. What is the percent yield of the reaction?

# Stoichiometry Example

What volume of H<sub>2</sub> gas (at STP) could be produced by the reaction of 1.00 kg of methane gas with excess water?



# Solution Stoichiometry

- Stoichiometry problems that involve solutions are just like other stoichiometry problems. **Molarity** serves as the *conversion factor* to get to and from moles.

- Example:**

What volume of a 0.250 M AgNO<sub>3</sub> solution is required to produce 5.00 g of AgCl when mixed with excess MgCl<sub>2</sub> solution?



## Solution Stoichiometry (cont.)

- If the problem asks for molarity, you will need to know both the moles of solute and the volume of solution.
- **Example:** What is the molarity of an  $\text{AlCl}_3$  solution if 100.0 mL of the solution completely reacts with 75.0 mL of a 0.110 M  $\text{Pb}(\text{NO}_3)_2$  solution?



# Stoichiometry Example

Consider the reaction of  $7.31 \times 10^{24}$  atoms of magnesium with 865 mL of 4.00 *M* hydrochloric acid (HCl). What is the theoretical yield (in grams) of H<sub>2</sub> gas?

