

TABLE 19.1

Standard Reduction Potentials at 25°C*

Half-Reaction	$E^\circ(V)$
$\text{F}_2(g) + 2e^- \rightarrow 2\text{F}^-(aq)$	+2.87
$\text{O}_3(g) + 2\text{H}^+(aq) + 2e^- \rightarrow \text{O}_2(g) + \text{H}_2\text{O}$	+2.07
$\text{Co}^{3+}(aq) + e^- \rightarrow \text{Co}^{2+}(aq)$	+1.82
$\text{H}_2\text{O}_2(aq) + 2\text{H}^+(aq) + 2e^- \rightarrow 2\text{H}_2\text{O}$	+1.77
$\text{PbO}_2(s) + 4\text{H}^+(aq) + \text{SO}_4^{2-}(aq) + 2e^- \rightarrow \text{PbSO}_4(s) + 2\text{H}_2\text{O}$	+1.70
$\text{Ce}^{4+}(aq) + e^- \rightarrow \text{Ce}^{3+}(aq)$	+1.61
$\text{MnO}_4^-(aq) + 8\text{H}^+(aq) + 5e^- \rightarrow \text{Mn}^{2+}(aq) + 4\text{H}_2\text{O}$	+1.51
$\text{Au}^{3+}(aq) + 3e^- \rightarrow \text{Au}(s)$	+1.50
$\text{Cl}_2(g) + 2e^- \rightarrow 2\text{Cl}^-(aq)$	+1.36
$\text{Cr}_2\text{O}_7^{2-}(aq) + 14\text{H}^+(aq) + 6e^- \rightarrow 2\text{Cr}^{3+}(aq) + 7\text{H}_2\text{O}$	+1.33
$\text{MnO}_2(s) + 4\text{H}^+(aq) + 2e^- \rightarrow \text{Mn}^{2+}(aq) + 2\text{H}_2\text{O}$	+1.23
$\text{O}_2(g) + 4\text{H}^+(aq) + 4e^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Br}_2(l) + 2e^- \rightarrow 2\text{Br}^-(aq)$	+1.07
$\text{NO}_3^-(aq) + 4\text{H}^+(aq) + 3e^- \rightarrow \text{NO}(g) + 2\text{H}_2\text{O}$	+0.96
$2\text{Hg}^{2+}(aq) + 2e^- \rightarrow \text{Hg}_2^{2+}(aq)$	+0.92
$\text{Hg}_2^{2+}(aq) + 2e^- \rightarrow 2\text{Hg}(l)$	+0.85
$\text{Ag}^+(aq) + e^- \rightarrow \text{Ag}(s)$	+0.80
$\text{Fe}^{3+}(aq) + e^- \rightarrow \text{Fe}^{2+}(aq)$	+0.77
$\text{O}_2(g) + 2\text{H}^+(aq) + 2e^- \rightarrow \text{H}_2\text{O}_2(aq)$	+0.68
$\text{MnO}_4^-(aq) + 2\text{H}_2\text{O} + 3e^- \rightarrow \text{MnO}_2(s) + 4\text{OH}^-(aq)$	+0.59
$\text{I}_2(s) + 2e^- \rightarrow 2\text{I}^-(aq)$	+0.53
$\text{O}_2(g) + 2\text{H}_2\text{O} + 4e^- \rightarrow 4\text{OH}^-(aq)$	+0.40
$\text{Cu}^{2+}(aq) + 2e^- \rightarrow \text{Cu}(s)$	+0.34
$\text{AgCl}(s) + e^- \rightarrow \text{Ag}(s) + \text{Cl}^-(aq)$	+0.22
$\text{SO}_4^{2-}(aq) + 4\text{H}^+(aq) + 2e^- \rightarrow \text{SO}_2(g) + 2\text{H}_2\text{O}$	+0.20
$\text{Cu}^{2+}(aq) + e^- \rightarrow \text{Cu}^+(aq)$	+0.15
$\text{Sn}^{4+}(aq) + 2e^- \rightarrow \text{Sn}^{2+}(aq)$	+0.13
$2\text{H}^+(aq) + 2e^- \rightarrow \text{H}_2(g)$	0.00
$\text{Pb}^{2+}(aq) + 2e^- \rightarrow \text{Pb}(s)$	-0.13
$\text{Sn}^{2+}(aq) + 2e^- \rightarrow \text{Sn}(s)$	-0.14
$\text{Ni}^{2+}(aq) + 2e^- \rightarrow \text{Ni}(s)$	-0.25
$\text{Co}^{2+}(aq) + 2e^- \rightarrow \text{Co}(s)$	-0.28
$\text{PbSO}_4(s) + 2e^- \rightarrow \text{Pb}(s) + \text{SO}_4^{2-}(aq)$	-0.31
$\text{Cd}^{2+}(aq) + 2e^- \rightarrow \text{Cd}(s)$	-0.40
$\text{Fe}^{2+}(aq) + 2e^- \rightarrow \text{Fe}(s)$	-0.44
$\text{Cr}^{3+}(aq) + 3e^- \rightarrow \text{Cr}(s)$	-0.74
$\text{Zn}^{2+}(aq) + 2e^- \rightarrow \text{Zn}(s)$	-0.76
$2\text{H}_2\text{O} + 2e^- \rightarrow \text{H}_2(g) + 2\text{OH}^-(aq)$	-0.83
$\text{Mn}^{2+}(aq) + 2e^- \rightarrow \text{Mn}(s)$	-1.18
$\text{Al}^{3+}(aq) + 3e^- \rightarrow \text{Al}(s)$	-1.66
$\text{Be}^{2+}(aq) + 2e^- \rightarrow \text{Be}(s)$	-1.85
$\text{Mg}^{2+}(aq) + 2e^- \rightarrow \text{Mg}(s)$	-2.37
$\text{Na}^+(aq) + e^- \rightarrow \text{Na}(s)$	-2.71
$\text{Ca}^{2+}(aq) + 2e^- \rightarrow \text{Ca}(s)$	-2.87
$\text{Sr}^{2+}(aq) + 2e^- \rightarrow \text{Sr}(s)$	-2.89
$\text{Ba}^{2+}(aq) + 2e^- \rightarrow \text{Ba}(s)$	-2.90
$\text{K}^+(aq) + e^- \rightarrow \text{K}(s)$	-2.93
$\text{Li}^+(aq) + e^- \rightarrow \text{Li}(s)$	-3.05

*For all half-reactions the concentration is 1 M for dissolved species and the pressure is 1 atm for gases. These are the standard-state values.